



A neutron star with a carbon atmosphere in the Cassiopeia A supernova remnant


Wynn C. G. Ho¹ & Craig O. Heinke²

2009/11/12

Kao, Kai-Wei

Institute of Astronomy, NTHU

An artist's impression of the
carbon-cloaked neutron star
Illustration: NASA/CXC/M.Weiss



Cassiopeia A is one of the youngest-known supernova remnants in the Milky Way and is at a distance of $d \doteq 3.4$ kiloparsecs (kpc) from the Earth.


- **Here they report an analysis of archival observations of the compact X-ray source in the centre of the Cassiopeia A supernova remnant. They show that a carbon atmosphere neutron star (with low magnetic field) produces a good fit to the spectrum.**

Chandra X-ray spectra of Cas A.

They considered archival Chandra X-ray Observatory data from two studies of Cas A, both using the ACIS-S charge-coupled device.

A series of Chandra observations, totalling 1 megasecond, was performed in 2004 to study the supernova remnant; these are referred to here as the Hwang data.

A shorter (70 kiloseconds) observation in 2006 was designed to study the compact source, here referred to as the Pavlov data.



WCG Ho & CO Heinke *Nature* **462**, 71-73 (2009) doi:10.1038/nature08525

Spectra from the Hwang (black) and Pavlov (red) observations and fits with their C spectral model. Error bars are 1 s.d. The lower panel shows the fit residuals, ΔS_{χ^2} in units of s.d.

- Previous works found poor fits to the data using magnetic ($B \geq 10^{12}$ G) H atmosphere spectra.
- Timing measurements suggest that they have (dipolar) magnetic fields of $B \ll 10^{12}$ G.
Neutron stars have weakly magnetic fields at birth. (Gotthelf et al. 2009)
- When $B = 2.35 \times 10^9 \times Z^2$ G ($\sim 8 \times 10^{10}$ G for carbon), magnetic-field effects on the radiation transport and atoms in the atmospheric plasma are negligible. (Rajagopal et al. 1996 ; Zavlin et al. 1996)
- If the magnetic field of Cas A is $\sim (1-5) \times 10^{11}$ G, then a spectral feature due to the electron cyclotron resonance may appear in the Chandra energy range, but this has yet to be detected. The lack of a visible pulsar wind nebula and no indication of magnetospheric activity like those seen in classical pulsars (with $B \geq 10^{12}$ G), also **suggest the magnetic field is low.**

Table 1 | X-ray spectral fitting of Cas A

Atmosphere model	N_H (10^{22} cm^{-2})	kT (eV)	Normalization	$\chi^2/\text{d.o.f.}$	Null hypothesis probability (%)
Hydrogen	$1.65^{+0.04}_{-0.05}$	241^{+7}_{-6}	$0.18^{+0.03}_{-0.03}$	106.3/99	29
Helium	$1.62^{+0.04}_{-0.05}$	228^{+9}_{-8}	$0.22^{+0.05}_{-0.04}$	112.4/99	17
Carbon	$1.73^{+0.04}_{-0.04}$	155^{+7}_{-6}	$1.84^{+0.56}_{-0.42}$	105.3/99	31
Nitrogen	1.37	172	1.18	388/99	0
Oxygen	1.03	234	0.20	2,439/99	0
Blackbody model	$1.46^{+0.04}_{-0.04}$	$kT^\infty = 387^{+7}_{-6} \text{ eV}$	$R^\infty = 1.0^{+0.1}_{-0.1} \text{ km}$	134.2/98	11

Joint fitting results to the Hwang and Pavlov data with neutron-star atmosphere and blackbody models that are modified by photoelectric absorption N_H (using the Tuebingen–Boulder absorption routine TBABS with Wilms model abundances in the X-ray spectral fitting package XSPEC; see <http://heasarc.gsfc.nasa.gov/docs/xanadu/xspec/index.html>), dust scattering²³, and corrections for pile-up²⁴ (the grade migration parameter of the pile-up algorithm is allowed to float freely and is 0.36 for the best-fit C model). Chandra data with such high statistics reveals systematic uncertainties in the Chandra calibration¹³, so we added a systematic uncertainty of 3% in quadrature. All parameter errors are given at 90% confidence. Errors are not reported when the reduced $\chi^2 > 2$. The normalization refers to the fraction of the neutron star surface emitting radiation (for a 10 km stellar radius and 3.4 kpc distance). The null hypothesis probability is the probability that one realization of the model fit to the data would have a reduced χ^2 greater than that obtained; less than 5% indicates a poor fit. $T^\infty = T/(1 + z_p)$ and $R^\infty = R(1 + z_p)$ are the temperature and radius measured by an observer at infinity and d.o.f. is the number of degrees of freedom.

$$\text{H} \rightarrow R \doteq 4 \text{ km}$$

$$\text{He} \rightarrow R \doteq 5 \text{ km}$$

$$\text{C} \rightarrow R \doteq 12 \sim 15 \text{ km}$$

$$R_{\text{NS}} \doteq 10 \text{ km} \sim 14 \text{ km.}$$


Thus we conclude that Cas A is consistent with a low-magnetic field carbon-atmosphere neutron star of mass $1.4M_{\text{Sun}}$, radius 12–15 km, and surface temperature $T=1.8 \times 10^6 \text{ K}$.

- The derived sizes of the emission region R for H and He are much smaller than the theoretical size of a neutron star R_{NS} . This would suggest the emission region is a hot spot on the neutron-star surface, which would probably result in X-ray pulsations as the hot spot rotated with the star. However, these pulsations have not been detected.

the X-ray observations are detecting emission from the entire stellar surface, and therefore the emission would not necessarily vary as the star rotates

Neutron star mass and radius.

Interpreting the size of the emission region to be the true neutron-star radius ($R=R_{\text{NS}}$), we can constrain the neutron-star mass M_{NS} and radius R_{NS} by using a range of surface gravity models. If we fix $M_{\text{NS}}=1.4M_{\text{Sun}}$ and the distance to Cas A as $d=3.4$ kpc, then we find a surface effective temperature of $T_{\text{eff}} \doteq 1.61 \times 10^6 \text{K}$ and emission size $R \doteq 15.6$ km.



WCG Ho & CO Heinke *Nature* **462**, 71-73 (2009) doi:10.1038/nature08525

90% confidence contours of χ^2 around the best-fitting Cas A mass and radius (crosses) for distances of 3.3 kpc (red long-dashed line), 3.4 kpc (thick black solid line), and 3.7 kpc (blue short-dashed line).

The upper-left and lower-right regions are excluded by constraints from the requirement of causality and from the fastest-rotating neutron star known.

- the compact source as a very young (330-year-old) neutron star.
- Because the next-youngest neutron stars for which surface thermal emission has been measured have ages exceeding a few thousand years, Cas A serves as a valuable window into the early life of a passively cooling neutron star.

• Thank you

• Reference:

Nature Vol 462|5 November 2009| doi:10.1038/nature08525

(<http://www.nature.com/nature/journal/v462/n7269/pdf/nature08525.pdf>)

Carbon atmosphere discovered on neutron star

(<http://www.astronomy.com/asy/default.aspx?c=a&id=8771>)